

DEPARTMENT of ENVIRONMENTAL SERVICES
Water Division - Watershed Management Bureau

LAKE TROPHIC DATA

MORPHOMETRIC:

Lake: CRANBERRY POND	Lake Area (ha):	7.49
Town: ALSTEAD	Maximum depth (m):	2.7
County: Cheshire	Mean depth (m):	0.9
River Basin: Connecticut	Volume (m ³):	64500
Latitude: 43°04'36" N	Relative depth:	0.9
Longitude: 72°17'14" W	Shore configuration:	1.86
Elevation (ft): 1135	Areal water load (m/yr):	25.21
Shore length (m): 1800	Flushing rate (yr ⁻¹):	29.20
Watershed area (ha): 391.2	P retention coeff.:	0.45
% watershed ponded: 0.9	Lake type:	artificial

<u>BIOLOGICAL:</u>		27 February 2002	31 July 2001
DOM. PHYTOPLANKTON (% TOTAL)	#1	DINOBRYON 70%	TETRAEDRON 90%
	#2	STAUSTRUM 20%	MOUGEOTIA 4%
	#3		EUDORINA 4%
PHYTOPLANKTON ABUNDANCE (units/mL)			
CHLOROPHYLL-A (µg/L)			5.59
DOM. ZOOPLANKTON (% TOTAL)	#1	SPARSE - NO DOMINANT	KERATELLA 27%
	#2		NAUPLIUS LARVA 21%
	#3		POLYARTHRA 14%
ROTIFERS/LITER		12	434
MICROCRUSTACEA/LITER		<1	285
ZOOPLANKTON ABUNDANCE (#/L)		14	761
VASCULAR PLANT ABUNDANCE			Abundant
SECCHI DISK TRANSPARENCY (m)			2.3 Visible on bottom
BOTTOM DISSOLVED OXYGEN (mg/L)		7.6	7.6
BACTERIA (E. coli, #/100 ml)	#1		< 1
	#2		< 1
	#3		

SUMMER THERMAL STRATIFICATION:

not stratified

Depth of thermocline (m): None
Hypolimnion volume (m³) : None
Anoxic volume (m³) : None

CHEMICAL:

Lake: CRANBERRY POND

Town: ALSTEAD

	27 February 2002		31 July 2001		
DEPTH (m)	1.5		1.5		
pH (units)	5.7		6.3		
A.N.C. (Alkalinity)	1.2		2.1		
NITRATE NITROGEN	0.05		< 0.05		
TOTAL KJELDAHL NITROGEN	0.20		0.50		
TOTAL PHOSPHORUS	<0.005		0.015		
CONDUCTIVITY (μ mhos/cm)	68.0		50.1		
APPARENT COLOR (cpu)	13		20		
MAGNESIUM			0.65		
CALCIUM			1.8		
SODIUM			5.5		
POTASSIUM			0.46		
CHLORIDE	13		9		
SULFATE	6		4		
TN : TP			33		
CALCITE SATURATION INDEX			4.4		

All results in mg/L unless indicated otherwise

TROPHIC CLASSIFICATION: 2001

D.O. S.D. PLANT CHL TOTAL CLASS


**	2	5	1	8	Meso.
----	---	---	---	---	-------

COMMENTS:

1. This pond is located within the Timber Owners of New England, Inc.'s Wildlife Conservation Trust and permission is required for access through the gate.
2. This is really two ponds – an upper and lower pond separated by a beaver dam.
3. Nineteen genera of net phytoplankton were observed, including many desmids, but there was a strong dominant and the dominant genus (*Tetraedron*) is one that is not generally dominant in NH ponds. Zooplankton were abundant and also varied (17 genera observed).

Cranberry Pond

Alstead



0 0.1 0.2 Kilometers

Cranberry Pond

Alstead



[illegible]